

7

Index
1 leaf to letter

FROM
THE LIBRARY
OF
SIR WILLIAM OSLER, BART.
OXFORD

7666 c

W. Osler Esq

University College

Toronto

Jan 1st / 1870

A

B

C

D

E

F

G

H

J

K

L

M

M'

N

O

P

R

S

T

U

V

W


Y

Z

notes in 1870
on animal Parasites etc.

A
B
C
D
E
F
G
H
J
K
L
M
M
N
O
P
R
S
T
U
V
W
Y
Z





B

C

D

E

F

G

H

J

K

L

M

M

N

O

P

R

S

T

U

V

W

Y

Z

Entozoa p. 175

- " " in dog. p. 200
" " , experiments with p. 230
" " in Bat 205
" " in Cat-fish p. 207
" " in Pig p 209
" " in Man p 211
" " in Chub p 217
" " " Pike p 219
" " " Perch p 223
" " " Eel p 227
" " " Squirrel p 229
" " " Pickerel p 250
" " " *Menobranthus lateralis* p 254
" " " Skunk p 258
" " " Sun-fish p 260
" " " King-fisher p 264
" " " Black Squirrel 266
" " " Bass p 268
" " " Hawk p 270
" " " Crane p 272

Entozoa in Fowl. p 274
 " " " Dony p 276 & 250
 " " " Bar-fish p 278
 " " " Cat p 280
 " " " Lynx p 282
 " " " Rat p 284
 " " " Beaver Fish (*Ameia calva*) p 286

E
 F
 G
 H
 J
 K
 L
 M
 N
 O
 P
 R
 S
 T
 U
 V
 W
 Y
 Z

Fish examined with negative results. p 453

F

G

H

J

K

L

M

M

N

O

P

R

S

T

U

V

W

Y

Z

Gatherings

- (Diatomace) p
- (Dermudica) p
- (Infusoria) p

G
H
J
K
L
M
M
N
O
P
R
S
T
U
V
W
Y
Z

H

J

K

L

M

N

O

P

R

S

T

U

V

W

Y

Z

Polyzoa p

P
R
S
T
U
V
W
Y
Z

20/3/70

I

Went out to Humber Bay with Rev W. A. S. and obtained
gatherings from Quader Pond and its outlet

24/3/70. Went out again and obtained another gathering
from the same places. The Diatoms were all alive and moving
freely. Many strips of *Ocellularia* were mingled with
them and gave the gathering a rather greenish appearance.
From the Pond on the right hand side of the bridge the following
Diatoms were obtained

Cymbella Ehrenbergii, *Cocconeis Shvaitzii*, *Pinnularia gibba*,
Navicula affinis, *Diatoma elongatum*, " " *distans*
Pinnularia major, *Stauroneis anceps*, *Cocconeis lance-*
Nitzschia sigmoides, " " *linearis*, *Navicula rhomboides*
Navicula lanceolata, *Navicula crassirostris*, *Epeithemia sorus*
Stauroneis punctata, *Savirella angusta*; *Cymbella cuspidata*
Navicula inflata, *Cocconeis cyrtiformis*, *Achnanthes exilis*
Cocconeis Aislula, *Gomphonema gemmatum*, *Gomph. acumin-*
Diatoma vulgare




a curious Rhizopod? outer line the clear protoplasm
the inner the boundary of the lorica



Erupted young Diatoms, probably *Cocconeis*,
they exhibit distinct movement in their cysts, and
then assume a vertical position



In this three cysts are seen



Navicula affinis, when ^{first} seen the endochrome was
rolled up into four balls, in a few minutes the
middle ones coalesced and enlarging pressed the
end ones against the apices. Seen with $\frac{1}{20}$

2/4/10 out at Weston, In a gathering from the ST Railway
numbers of *Nitzschia amphioxys* and *Mendion curv*
are

Ernadestond 23/3/70

- Cymbella Ehrubergii*
- Cocconeis thwaitesii*
- Pinnularia gelba*
- Nauicula affinis*
- Diatoma elongatum*
- Pinnularia ^{distans} ~~peruviana~~*
- " " *major*
- Stauroneis anceps*
- Cocconeis lanceolata*
- Nitzschia sigmoides*
- Stauroneis linearis*
- Nauicula rhomboides*
- " *lanceolata*
- " *crassirostris*
- Ephemia corex*
- Stauroneis punctata*
- Sarivella angusta* 1/20
- Hemidictyon of rudochromis*
- Pinnularia gelba* 1/20



u. l. m. m. s. 1/20
 The upper one would be the clear setae
 in the middle of the
 in the middle of the
 in the middle of the



The upper one would be the clear setae
 in the middle of the
 in the middle of the
 in the middle of the



in this one there are
 in this one there are
 in this one there are
 in this one there are

The rudochromis in this is rolled
 up into four
 the largest
 two center
 spread them
 through out the cell
 of passing the smaller ones
 along the apices. This was in *Nauicula affinis*
 and seen with 1/20 2/2/70



Herbert Pond 23/3/70

Cymbella curvata

Nannula inflata

Cocconeis cymbiforme

Achnanthes exilis

Cocconeis cistula

Gomphonema geminatum

" " *accumulatum*

Diatoms vulgares

In this gathering from the Herbert Pond
several infusoria coloured an ochre
brown, probably from the eating the brown
coloured fungus *Cothurnia*

W

Entozoa

Entozoa in Dog

19/2/69

I

From a ^{young} slut; three or four *Ienia elliptica*, two *Bothriocephalus cordatus* (slide) and a number of ascarida. This animal had been about a butcher's shop

II

22/1/70

From a slut, six months old, also got from a butcher's shop. 39 *Ienia elliptica*, 25 *Ascaris*. In this dog, there was but one species of *Ienia*, the *Ienia* are collected towards the ileum, while the ascarida are usually in the duodenum & beginning of jejunum. Of the 39 *Ienia* I obtained 18 with heads, the other came away without, breaking off close to the head. Besides the strobila, there were numberless mature ^{prep} proglottides; the eggs contained in these exhibited distinctly the six-hooked embryo = proscelices. Slides No's , of heads; No's , of ova

7/2/70 No III
 From a ^{temer} slut about a year old, four ascaridæ.
 No *Tania* at all

IV.

4/3/70 From a dog about a year old, which had on a former occasion 1/3/70 been fed with flesh from Case IV p 212.
 4 *Ascaris* and four *Tania elliptica* all with heads were obtained. No *Inchima* were found either in the intestines or in the muscles, but on examining the kidneys. 12 or 13 very small white spots were seen about $\frac{1}{2}$ that size (see slide) in each of these ~~was~~ seen a small *Normalode* worm was observed not coiled up, and looking exceedingly like a *Trichina spiralis*

V

14/2/70 From a pup about four months old, which had previously been fed with *Inchima*. 3 *Ascaris* were obtained

14/2/70 Examined a fine St-Bernard dog. No entozoa

10. 12. 19. 20. 21. 22. 23. 24. 25. 26. 27. 28. 29. 30. 31. 32. 33. 34. 35. 36. 37. 38. 39. 40. 41. 42. 43. 44. 45. 46. 47. 48. 49. 50. 51. 52. 53. 54. 55. 56. 57. 58. 59. 60. 61. 62. 63. 64. 65. 66. 67. 68. 69. 70. 71. 72. 73. 74. 75. 76. 77. 78. 79. 80. 81. 82. 83. 84. 85. 86. 87. 88. 89. 90. 91. 92. 93. 94. 95. 96. 97. 98. 99. 100.

Entozoa in Bat

15/11/69

obtained a curious encysted Nematode from
the walls of the stomach of the New-york Bat

, there were no other Entozoa, either free in
stomach or intestines or encysted in the muscles

There were eight altogether, Slides no , see drawing

Protozoa in Cat-Fish

12/5/69

Procured two small *Tania* from intestine, length about half an inch; head devoid of hooklets

7/6/70 From a cat-fish caught in the canal 10 *Tænia* were obtained. This cestode is from $\frac{1}{4}$ to $\frac{1}{2}$ an inch in length and very extensible. Suctional disks prominent; can be seen with the naked eye they are four in number; no hooklets were seen. The water vascular system is beautifully seen in this species at the head it may be seen terminating in a plexus (whose branches are as large as the primary trunks) situated between the disks

Two *D. stromes* were also found in the intestine
 Encysted Nematodes in the liver
 a *Cyrtocercus* in the Kidney

29/5/70

From the intestines of Cat-fish speared at the Island. 3 *Tania* very small and 4 Echinorhynchide (see slide at W. a. S.). accessory claspers of the male well seen in

Entozoa in Pig

5/6/69

Obtained from Dr Maddock of Hamilton a small portion of the heart of a measeley pig, contained about a dozen of *Cyrtocercus Cellulosa*. See Slides No

Entozoa in Man

7/2/68

Trichina spiralis

I

Obtained a piece of muscle from a man who died of the disease in Illinois State, very numerous not encysted, Two families living on different flats of the same house bought a barrel of Pork between them, one flat cooked the pork and escaped the disease, the other eat their raw and nine were sickened; of these nine, four died

27/2/69

II

The family of a Mr Setz in Hamilton consisting of himself, wife & daughter, put pork of an uncorked ham all three were laid up with the disease, Mrs Setz died first, in her the parasites were numerous & unencysted, Mrs Setz died some two weeks after her daughter in her they were just beginning to be encysted. The husband was attacked but not so severely and escaped, most probably from being drunk for some days at the commencement of the attack, See Slides No's

20/2/69

III

Obtained a piece of muscle from a woman

in New York. In her the disease was not suspected during life, the students on dissecting the body found (as in the original discovery of the parasite) great difficulty in cleaning the muscles, and on examination they were found to be packed with Trichinae, all encysted. See Index No's

22/2/70

IV

While dissecting the arm of a man, who had died at the Toronto General Hospital, I found numerous Trichinae in the Biceps muscle and further examination showed them to be scattered pretty freely throughout the muscle. From 31 of the muscle from the beginning of the long head of Biceps I obtained 150 cysts, the greater number of them containing healthy looking Trichinae

29/3/70

V

In the subject following the one above, and ^{also} brought from the Toronto General Hospital, numerous cysts were found in all parts of the body. The parasites in this case are not as old, none of them beginning to undergo degeneration.

Entozoa in Man

Taenia solium 15/3/68

Im pet passed by a bank-clerk
a patient of Dr. Russell's, see slides No's

2 3/2/71

Taenia mediocanellata. Obtained three specimens

of this cestode from a man ^(Earle) who died of heart disease
in Montreal Gen Hospital. He stated during life that

he had been suffering from tape worm for 14 years
got it at Matta while stationed there with his regiment.
He stated that he had been under treatment for it
innumerable times & had passed many yards of the worm
while in Hospital during the end of last year & the
beginning of this he was treated with the male shield
fern, which brought away several long portions.

On opening the small intestines the worms were found
extending from the lower part of the duodenum, thro-
ugh the jejunum & seeming completely to fill the intestine
in the empty condition. The heads were all within 1/2 inch
of each other & deeply impeded in the mucous mem-

trane, between the valvulae conniventes. The bodies were
 convoluted & twisted, extending down the intestine for
 about $3\frac{1}{2}$ feet, & lower down in the bowel several detached
 portions were found consisting of from sixteen to
 eighteen proglottides. The worms still retained some
 little vitality, but the movements were very feeble. While
 in the intestine, the water vascular system in one of the
 worms beautifully seen, extending up each side of the
 worm. The length was of each respectively 76.50 &
 65 inches & the number of segments enumerated in
 each to between 275 & 350, falling at least 100 short
 in the longest one of the fully mature sexual seg-
 ments. This however is not to be wondered at as this not
 more than six weeks ago since he passed several yards.
 The chief differences between this form & *I. solium* apper-
 tly as follows. The head is larger, abruptly truncated
 & lacks the scolextum & consequently the hooklets. This
 surrounded by dark zone of calcareous corpuscles which
 form a striking contrast to the white segments of the
 neck. The segments are broader, thicker & not as long.
 The generation orifice is a little below the center

& the lateral branches of the uterus, appear more numerous & closely packed together. There also appears to be a difference in the shape of the ova in the two species that in *Icenia solum* being round, while in *I. marginata* they are rather oval & somewhat larger.

On the fins of chub

2/4/70 In the Rev W. a. Johnson's aquarium, were noticed several round white spots, here on examination proved to be some sort of Intozoa. In addition to these ~~white~~ some yellow spots were seen which seem to be a more advanced condition of the parasite (see slide)

2/5/70 Numerous flukes attached to the intestine of a small chub. (see slide no)

12/6/70 Examined 3 chub. from the intestine of one two *Echinorhynchus* were obtained, a male & female

Entozoa in Pike

23/4/70

In a Pike 2 ft 7 inch long caught at the Island I obtained 68 specimens of *Tania* and two or three small ascarids

This tape-worm is about a foot long, ~~very~~ exhibited curious undulatory movements which continued for more than twenty four hours after removal from the intestines. It is very extensible and may be stretched to almost double its ordinary length. The head is flattened, club shaped when the worm is dead but during life is generally extended giving to ~~the~~ it the shape of a flat arrow head. ^{7 side} Four suckorial disks are plainly seen but no ~~hooklets~~. The segments taper very gradually being exceedingly small at the neck, larger towards the end of the body, they are about twice as broad as they are long. The water vascular system is not distinctly seen in this worm consisting of ^{four} two channels ^{two} on each side, at the head and for a considerable distance down the neck these tubes connect by means of vasculating branches, these about the head form a dense net-work (see sketch)

30/4/70 From a pike caught in the canal basin at Dundas, I obtained 28 Tania and numerous small Ascaridae. In the stomach of this fish were 52 smaller ones, principally little bass perch

3/6/70

From the intestines of two Pike obtained at the Fish-market Toronto. In one 84 Tania were found and in the other 53 not counting numerous small undeveloped ones, looked like freshly eaten scolices. A few Ascaridae were found in the stomach of one.

23/6/70 From intestine of a Pike obtained in Fish-market Toronto. 56 Tania; most of these were of a large size and longer than the usual ones from this fish

Perch
Entozoa in ~~Chab~~

23/4/70 In the liver of a perch taken from the stomach of a Pike (see p. 219) were numerous encysted nematodes (see slide no)

13/5/70 From four Perch caught in the Canal I obtained the following.

A Cyrtocercus from the liver, the head of which bears a remarkable resemblance to the head of the same of the Pike & most probably is its scolex.


Encysted Nematodes from the liver

Four species of Echinorhynchus from the rectum


In the intestine there was a curious Entozoon with a bell-shaped retractile head furnished with small finger like processes. (see sketch slide)

Coiled around the base of the aorta was a large Distome the upper tierid is outlined in this sketch

9/5/70



From the duodenum of eel, one large worm
and two of they were from $\frac{3}{4}$ to 1 inch
in length, moved freely, retracting and pushing
out the hooked proboscis, hooklets 11 deep, &
5 or 6 in each row, curved like the bars of a
harpoon, the body was filled with ova, enclosed
in a tube extending the whole length of the
worm. Two constrictions great near the
head but none in the rest of the body



225

MCGILL UNIVERSITY
MONTREAL

MEMO FROM
MEDICAL LIBRARY

TO:

Transfer to Sec. 4.

Older Role please

Chris

DATE _____

Entozoa in Sel

9/5/70 one solitary tape-worm in the duodenum about four inches long, no generative organs seen, no ova, water-vascular system distinct; no head discovered

Two species of Echinorhynchus both females length $\frac{3}{4}$ to one inch, moved freely, retracting and pushing out the hooked proboscis, eleven rows of hooklets, recurved like the barb of a harpoon the two Lemnisci were distinct - as also the three muscular retractors of the proboscis. Cysts containing ova occupied the rest of the body. Two or three slight constrictions noted

12/7/70 From a large silver eel caught at Burlington canal. Obtained one large DuLome from stomach

229

Entozoa in Squirrel

12/5/70 Eight small *Oscarida* from the duodenum
of the red Squirrel

23/1/70

Experiments with Entozoa

†

Fed two rabbits with mature Proglottides of *Tenia elliptica*, from Dog no 2, Six segments to each rabbit
 These will in all probability give negative results as the scolices of *Tenia elliptica* are not produced in the rabbit's
 These failed.

††

23/2/70

Fed a rabbit with about 3 ⁱⁱⁱ of *Inchinos* flesh from man in whom I found them while dissecting

1/3/70

†††

Fed a cat and dog with *Inchinos* flesh 3/3/70 repeated the dose to dog


In these cases the juices of the stomach were in all probability not strong enough to dissolve the cyst and free the parasites

 IV


30/3/70 Fed a rabbit and a young pup with muscle from
 Case I. 31st repeated the dose to pups. 1st again.

14/4/70 Killed the pup. No traces of Trichinae; the cyst was
 probably too dense for the juices of the stomach to dissolve.

21/4/70 Killed the rabbit fed with the Trichinosis flaps
 on 30th March. Numerous Trichinae in a young
 immature condition were observed in the muscles.
 Many of them exhibited sluggish movements,
 they were more numerous in the abdominal and
 thigh muscles than in any other.




13/5/70




canine
 + 1. inch

Head of perch from intestine
 see Chile




18970

Head of *Tania* from S. Kruak
Camera Lucida. and high object-



245

247



Entozoa in Pickeral

3/6/70

From the intestine of a Pickeral in the Fish market I obtained 8 Tania. This is a comparatively short cestode length from 10-12 inches, breadth of posterior segments about $\frac{1}{4}$ of an inch. Head square, no neck but a gradual tapering to the head, suckers four, large, no proglottids observed.

23/6/70 From Pickeral (1 foot - the half long) in Fish market Toronto. 18 Large Tania were obtained.

5-15-1908

1000

[Faint, illegible handwriting on lined paper]

253

Entozoa in *Meurobranchus lateralis*

6/6/70

From a specimen of above, caught at the Island I obtained numerous Polystomes, some of which were attached to the branchia others to the upper surface of the mouth. This trematode presents four disks, two at each extremity. Of these the smaller one situated about the middle of the upper fourth of the body, is the mouth, which leads directly into the intestines, ^{these} consisting of two simple tubes, which unite about the middle of the lower fourth of the body. The water vascular system is well developed. It appears to commence in a ramification of vessels about the anterior disk, these unite to form two vessels, which run the whole length of the body, join below and open somewhere between the posterior disks. Cilia as to be seen distinctly in the water vascular system, especially at the junction of the tubes below. At the ~~far~~ upper third of the body on a level with the generative opening, on each side is seen a curious pulsating organ, which is undoubtedly connected with the water vascular system. The pulsation occurs about once every minute & half. The external generative orifice is seen a little below the

the oral disk, this leads into a narrow slightly curved vagina. The ova occupy the general cavity of the body between the digestive tubes, a bundle of spicules (Pennis?) are to be seen close to the vulva. Close to the caudal disk two large hooks are placed. These hooks are bifurcate and to one extremity a band of fibres is attached. Besides these numerous small hooklets are to be seen scattered down the posterior disk. The attached extremity of these is trifurcate.

257

Entozoa in Skunk

15/6/70

From a large male Skunk. About 30 *Taenia* and 14 *Ascarida*. Numerous small cysts were ^{observed} ~~found~~ in the liver & spleen, but nothing found in them. The Tape worms are small from $\frac{1}{2}$ an inch to 2 inches ^{in length} broad in proportion to their length, exhibiting very slight movements. A slight enlargement seems to exist about the neck, which disappears when the worms is much elongated. The segments seem but loosely joined together breaking very easily. Four larger sucking disks exist at the head, no sucklets seen. The calcareous corpuscles are more numerous in this cestode than in any I have yet examined. The water-vascular system is not easily seen on account of the dense layer of Calc. corpuscles.

The *Ascarida* are from $\frac{3}{4}$ - 1 inch in length. They move freely. Most of them were in the stomach not in the intestines.

259

Journal in 1871

Intozoa in Sun-fish


260

4/7/70

Examined 10 Sun-fish caught in the Canal; in all numerous Distomes were found. This fluke is probably in an immature condition, being encysted and not having its internal organs completely developed. The heart, liver & Kidneys presented a mottled appearance from the numbers in them. They seemed only to be attached to the heart, while in the liver & Kidneys they occupied the substance of those organs,

Encysted Nematodes were found in one liver and what appeared to me the scolex of one of the Echinorhynchida, in another.

In the rectum of two, a few Echinorhynchida were found





263

Entozoa in King-fishes

264

17/70

Examined one of these birds and much to my surprise found no Entozoa in it

13/8/70

Shot a Kingfisher a few small Distomes were found in the liver. The small fish which constitute the food of this bird seem not to share the common fate of fish, in as much as few mature or immature Entozoons are found in them

265

Entozoa in Black-Squirrel

1/7/70

Examined a large male animal. No Entozoa
in them

267

Entozoa in Bass

7/7/70

Examined a large Black Bass caught in Burlington Bay, a solitary scolex of some tape worm was found in the peritoneal cavity

Intozoa in Hawk

13/8/70

Shot a Hawk; a single large Ascaris found
in the duodenum

Intozoa in Crane

15/8/70

Entozoa in Fowl (*Gallus ~~Bankiva~~*)

9/9/70 Examined the intestines of three fowls, one a two year old Rooster the others, this year's chickens. In the duodenum of the Rooster numerous *Tania*, from 2-6 inches long, were found; in the younger fowls a few smaller ones of the same kind occurred. The smaller ones were deeply imbedded in the mucous membrane the larger ones not at all so.

4/1/71

Examined intestines of two fowls, found in one numerous *Tania* same kind as above, and one *Distome* about half an inch long & 2 lines thick, very large posterior sucking disk. The fowl has been frozen so that the worms were dead & had begun to disintegrate. In the other a few small *Tania* only were found. The generative organs in this species open by an orifice of considerable size along one margin only of the worm. The opening is more at the junction of the segments than at the center.

6/1/71 Examined two fowls, one large ascaris only found. The fowls were grumped.

Intozoa in Dory

22/10/70

Examined a dory in Montreal fish market. One small tape worm, one Bothnocephalus and about two dozen small Echinorynchidae were found. The Bothnocephalus occupied the enter cavity of one of the numerous ceca given off at the pylorus, the Tænia was in the duodenum & the Echinorynchidae more towards the rectum. This fish I find in the Pickens
 see p 260

Entozoa in Bar-fish

4/11/70 Examined one of the above named fish, and found in the peritoneal cavity some half dozen Nematode worms. Each worm is about half a line in thickness and from three to four inches in length, dark in colour, annulated towards the anterior extremity, which is blunted and thick, the posterior extremity is pointed & presents the vaginal & rectal orifices. The ovaries extend the whole length of the body.

10/11/70

Found more of the same kind in the peritoneal cavity, passed about the rectum, nothing in the intestines.

Entozoa in Cat

280

3/1/71

Examined intestines of cat, found one *Ascumyctax* (?) and one *Tænia*

Eutozoya in Lynx

282


5/3/71 Examined the intestines of an animal obtained by the Nat History Society, Montreal. In the duodenum were ascandae & lower down in the bowel 25-
Janice. The animal had been dead a long time & the intestines had been thrown out & exposed to the cold for two nights. The greater part of a racoon was found in a semi-digested state in the stomach. The other organs were not examined

Eutoyoa in Rat

284

8/3/71

From a rat obtained at Montreal General Hospital
I obtained 5 Tamea. They were situated low down in the intestines,
approaching the faeculent matter. This is a small, fine
species. $2\frac{1}{2}$ to 3 inches in length. Head very small no hooks seen.
The water vascular system is very distinct.



Tapeworm from intestine of Beaver Fish, (*Amia calva*),
magnified.
A. H. Foord, del.

451

Fish examined with negative results

4/4/70

Examined two small Sturgeons, contents of stomach, numerous larva of all kinds, principally those of the Libellula
 Examined a large Sucker, stomach contained small water corns
 and larva

10/11/70. a woodrized Sturgeon, two Muschelunge, a mullet & a
 Pickarel

